


IMMUNOMODULATION WITH SYNTHETIC MOLECULES: MECHANISMS  
OF ACTIONS AND EFFECTS ON MACROPHAGES(U) INSTITUTE FOR  
MEDICAL RESEARCH SAN JOSE CA D A STEVENS ET AL.

NL

01 NOV 84 N00014-83-K-0018

F/G 6/5

END



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963 A

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

2

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 3	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) IMMUNOMODULATION WITH SYNTHETIC MOLECULES: Mechanisms of actions and effects on macrophages		5. TYPE OF REPORT & PERIOD COVERED Annual Report 83NOV01 to 84OCT31
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) David A. Stevens, M.D., Elmer Brummer, Ph.D, Christine J. Morrison, Ph.D, Dept. of Medicine, Stanford Univ. Med. School, Stanford, CA		8. CONTRACT OR GRANT NUMBER(s) NU0014-83-K-0018
9. PERFORMING ORGANIZATION NAME AND ADDRESS Institute for Medical Research of Santa Clara County, 2260 Clove Drive, San Jose, CA 95128		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 666-013
11. CONTROLLING OFFICE NAME AND ADDRESS Jeannine A. Majde, Ph.D., Scientific Officer Immunology, code 441, Cellular Biosystems Grp., Dept of the Navy, ONR, Arlington, VA 22217		12. REPORT DATE 34NOV01
		13. NUMBER OF PAGES 4
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Same as 11		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Unlimited <b>DISTRIBUTION STATEMENT A</b> Approved for public release; Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Unlimited <b>DISTRIBUTION STATEMENT A</b> Approved for public release; Distribution Unlimited		
18. SUPPLEMENTARY NOTES B		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Immunology, immunomodulators, immunoadjuvants, immunoenhancers, fungi, fungal infection, mycoses, host defenses, glycopeptides, muramyl dipeptide, lymphocytes, macrophages, polymorphonuclear leukocytes, microbial killing, lymphokines, lipoidal amines, interferon, interleukin-1, interleukin-2, alveolar macrophages, phagocyte oxidative metabolism, chemiluminescence		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Immunoenhancing drugs are under study both for their general effects on host immunity and their effects on immune interactions with microbial infections. These studies focused preferentially on defined synthetic single molecules, especially synthesized natural biological molecules or their active fragments. We studied the mechanism of muramyl dipeptide in inducing hyporesponsiveness of spleen cells and hyperresponsiveness of lymph node cells to mitogens defining		

DD FORM 1473

1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

S/N 0102: LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

84 12 04 088

AD-A148 592

DTIC FILE COPY

DTIC

ELECTE

DEC 13 1984

B

the cell populations and mediators involved in these effects.

In another study, murine macrophages from different anatomic compartments were compared for their fungicidal activity in different in vitro systems. Subsequently, the effect of a lipoidal amine immunomodulator on alveolar macrophage fungicidal activity was examined.

The mechanism of immunological activation of polymorphonuclear neutrophils for fungicidal activity was studied with respect to the oxidative mechanisms involved.

Recombinant gamma interferon was shown to activate macrophages for fungal killing.



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Our work on immunomodulation and its mechanisms continues. The principal efforts of the past year, work in progress and future efforts are described below.

## I

Earlier in the year we concluded the second phase of our studies on the mechanism of immunomodulation by muramyl dipeptide (MDP). We previously reported that a MDP treatment regimen (200  $\mu$ g MDP per mouse, days -4, -3, -2, -1) that affords protection given prophylactically against several infectious agents also induces lymph node hyperplasia, and lymph node cell (LNC) hyperresponsiveness to mitogens, whereas spleen cells make depressed responses to mitogens. The purpose of the present work was to extend those studies and delineate cellular mechanisms involved in these phenomena. It was found that hyperresponsiveness of LNC was prolonged (7 days) post treatment; in contrast hyporesponsiveness of spleen cells was transient and rebounded by day 4 post treatment. Hyperresponsiveness of LNC and hyporesponsiveness of spleen cells actively enhanced or depressed normal lymphoid cell responses, respectively, in cell mixing experiments. Hyporesponsiveness of spleen cells was associated with the plastic nonadherent, non-B-cell fraction and nylon wool nonadherent subpopulations. Indomethacin ( $10^{-6}$ M) did not abrogate hyporesponsiveness of spleen cells. These data suggest that splenic suppressor T-cells result from MDP treatment and were responsible for spleen cell hyporesponsiveness. On the other hand, hyperresponsiveness of LNC was associated with the nylon wool adherent cell subpopulations and a higher percentage of nonspecific esterase positive cells. Hyporesponsiveness of spleen cells was associated with deficient production of interleukin-2 (IL-2), but not of interleukin-1 (IL-1). In contrast, hyperresponsiveness of LNC was not explained by enhanced IL-1 or IL-2 production.

## II

Because of a need to extend our studies of effector cell function to alveolar macrophages, we performed further studies of alveolar macrophages in our killing systems with fungal targets. These studies extend our observations previously reported with these cells and Candida albicans (Amer. Rev. Resp. Dis. 127:110-112, 1983)

Murine bronchoalveolar macrophage (BAM) fungicidal activity against the yeast form of a virulent strain of Blastomyces dermatitidis was studied in the small volume wells of a Terasaki tray. In a four hour co-culture assay, significant killing (16-33%) of the fungus was demonstrated with resident BAM and with BAM obtained from mice rendered immune to lethal infection with B. dermatitidis. No significant differences between the activity of BAM from these two sources could be identified. Addition of 10% autologous normal or immune serum did not augment macrophage fungicidal activity. Simultaneous experiments with peritoneal macrophages (PM) also gave reproducible killing of the yeast form in wells of the Terasaki tray. However, PM studied in a standard microtiter plate assay showed no statistically significant fungicidal activity. On the other hand, BAM had similar fungicidal activity against B. dermatitidis in Terasaki and microtest plate wells. Modest fungicidal ability of resident BAM in immune mice against B. dermatitidis suggests that the striking resistance of immune mice to respiratory challenge is likely based on some augmentation of this first line of defense.

### III

Our studies of the lipoidal amine synthetic immunomodulator CP46665 also continued. Our results with polymorphonuclear neutrophils (PMN) were reported previously. Further studies with BAM were performed, which failed to demonstrate a significant effect, not confirming our earlier promising results. These experiments were done at the apparent optimal dose of 5 mg/kg, in a single dose, 1 day prior to experiment, using Terasaki plates. The control is the diluent for the intravenous injection, intralipid.

TABLE 1  
PERCENT KILLING OF CANDIDA

<u>Experiment #</u>	<u>Intralipid</u>	<u>CP46665</u>
1	10	43
2	17	47
3	18	18
4	22	20
5*	14	5
6	12	23

\*Utilized log phase *C. albicans*, others used plateau phase

If we omit experiment 5, which was the only one not identical in technique to others, the data can be analyzed as follows. The means  $\pm$  S.D. are  $16 \pm 5\%$  for IL and  $30 \pm 14\%$  for CP, which (apparently because of the large CP S.D.) is  $p > 0.05$ .

We also performed experiments when the dose was given in 2 consecutive days prior to the experiment. These 1 dose and 2 dose regimens, with sampling the effect on cells the following day, appeared optimal in early dose-finding experiments.

TABLE 2  
PERCENT KILLING OF CANDIDA

<u>Experiment #</u>	<u>IL</u>	<u>CP</u>
7	5	11
8	3	15
9	6	3
10	10	15

Applying the same analysis, the means  $\pm$  S.D. are  $6 \pm 3\%$  and  $11 \pm 6\%$ . Thus, even though 3 of 4 experiments show an edge for CP, the  $p$  is still  $> 0.05$ .

### IV

Recent studies have focused on the mechanism of the interaction between the T-cell mediated immune system and polymorphonuclear leukocytes (PMN) we previously described. We have shown, as previously described, using fungi (three different genera) as the target organisms, that the PMN can be modulated by lymphokines to new or enhanced levels of killing ability.

PMN induced locally in immune mice by intraperitoneal injection of antigen exhibit enhanced fungicidal activity compared to PMN elicited with thioglycollate. The mechanism of the differences in these PMN populations was studied. Sublethal infection was used to produce immunity to B. dermatitidis. A correlation was sought between ability of PMN to kill (or not kill) B. dermatitidis and production of the oxidative burst, as measured by luminol-enhanced chemiluminescence. Although elicited PMN cocultured with C. albicans produced a burst of chemiluminescence and were candidacidal, killing did not occur when the PMN were co-cultured with B. dermatitidis. Lack of killing of B. dermatitidis by elicited PMN correlated with lack of stimulation of a brisk oxidative burst. In contrast to elicited PMN, PMN induced by B. dermatitidis antigen responded to this fungus with a burst of chemiluminescence and a significant reduction of inoculum colony forming units (80%). Furthermore, these PMN when co-cultured with C. albicans produced an enhanced burst of chemiluminescence, and killing was enhanced compared to elicited PMN, e.g. 86 vs. 53%. The chemiluminescence burst and killing of B. dermatitidis by antigen-induced PMN was abrogated in the presence of catalase, implying a critical role for hydrogen peroxide. Partial but significant depression of chemiluminescence and killing in the presence of dimethyl sulfoxide, a hydroxyl radical scavenger, identified hydroxyl radical, or its metabolites, as toxic product(s) responsible for a significant fraction of fungicidal activity. These results indicate that the metabolic activity and microbicidal activity of PMN can be altered (enhanced) at the site of an immunological reaction, and thus could constitute an important factor in resistance.

#### V

Although our earlier reported studies indicated no beneficial effect of recombinant alpha-interferon on macrophages with respect to fungal targets, and in some instances a deleterious effect (these studies included human IF, C. albicans, human PMN and monocytes, and phagocytosis and killing), further studies with gamma interferon were indicated.

Recombinant murine gamma-interferon (IF) and supernatants from concanavalin A (ConA) stimulated mouse spleen cells (ConA-Sup) were tested for their ability to activate resident murine PM (peritoneal macrophages) (90% esterase positive) for fungicidal activity. PM monolayers pulsed overnight with IF had significantly enhanced fungicidal activity against C. albicans. ( $44 \pm 12\%$  vs.  $0.0\%$ , mean  $\pm$  SD of 3 experiments) and B. dermatitidis ( $34 \pm 1\%$  vs.  $3 \pm 3\%$ ). The effect was dose dependent; however, less IF ( $10 \text{ U/ml}$ ) was required to optimally activate PM to kill phagocytizable C. albicans than for nonphagocytizable B. dermatitidis ( $10,000 \text{ U/ml}$ ). ConA alone ( $5\text{--}10 \text{ mcg/ml}$ ) enabled PM to kill C. albicans but not B. dermatitidis. ConA-Sup activated PM for fungicidal activity against B. dermatitidis. ConA-Sup collected at 24 hrs. best activated PM for killing ( $30 \pm 3\%$ ) compared to those at 48 or 72 hrs. Capacity to activate PM for killing C. albicans or B. dermatitidis was abrogated when IF was incubated with anti-mouse IF monoclonal antibody. C. albicans killing was still induced after neutralizing ConA in ConA-Sup by absorption with Sephadex G-10 or incubation with alpha-methyl mannoside. Lipopolysaccharide ( $0.001\text{--}10 \text{ mcg/ml}$ ) failed to consistently activate PM for fungal killing. These data show that IF can have an immunoregulatory role in macrophage defense against these fungal pathogens, and may have therapeutic potential.

Future studies planned. With respect to section I above, these may include (a) further definition of the cell subsets affected by MDP, and (b) determining the mechanism of MDP protection against Blastomyces infection in vivo, by assaying humoral immunity and cell-mediated immunity (including macrophage) function in our model after in vivo MDP administration. With respect to section III, we will be completing some in vivo studies with CP 46665. With respect to section IV, the studies will progress in the following way. One avenue of research will pursue the biochemical mechanisms in the PMN consequent to immune activation of PMN's via lymphokines that result in

increased or new killing power. The second will be to characterize the lymphokine involved. The third avenue will attempt to define biological modifiers, produced by recombinant DNA or hybridoma techniques, which may produce the same effect. Prime candidates are gamma interferon and interleukin-2. If this is uncovered, it could lead to a therapeutic approach to defense against fungal pathogens. The last avenue will be to explore the effect of other eliciting agents, which may cause release in vivo of inflammatory mediators that could result in enhanced killing compared to the results with thioglycollate, and metabolic effects and killing power comparable to elicitation in immune animals with antigen, as we have described. With respect to section V, we are interested in the biochemical mechanisms in the macrophage that correlate with the enhanced fungicidal activity following gamma interferon treatment.

#### Publications

1. Brummer, E., Sugar, A.M., Stevens, D.A. Immunological activation of polymorphonuclear neutrophils for fungal killing: studies with murine cells and *Blastomyces dermatitidis* in vitro. *J. Leukocyte Biol. (RES)* 36:505-520, 1984
2. Brummer, E., Stevens, D.A. Activation of murine polymorphonuclear neutrophils for fungicidal activity with supernatants from antigen stimulated immune spleen cell cultures. *Infect. Immun.* 45:447-452, 1984
3. Williams, P.L., Brummer, E., Stevens, D.A. Effect of human prealbumin fraction with thymic hormone activity on the mononuclear cells of coccidioidomycosis patients and tuberculosis patients. In, *Coccidioidomycosis: Proceedings of the 4th International Symposium on Coccidioidomycosis*; H. Einstein, A. Catanzaro, eds.; Natl. Foundation for Infectious Diseases, pub; in press.
4. Brummer, E., Beaman, L., Stevens, D.A. Killing of endospores, but not arthroconidia, of *Coccidioides immitis* by immunologically activated polymorphonuclear neutrophils. In, *Coccidioidomycosis: Proceedings of the 4th International Symposium on Coccidioidomycosis*; H. Einstein, A. Catanzaro, eds; Natl. Foundation for Infectious Diseases, pub; in press.



**Annual, Final and Technical Reports (one copy each except as noted)**

Dr. Jeannine A. Majde, Code 441CB  
Scientific Officer, Immunology Program  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217

Administrator (2 copies)  
Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314

**Annual and Final Reports Only (one copy each)**

Commanding Officer  
Naval Medical Command  
Washington, DC 20372

Commanding Officer  
Naval Medical Research & Development Command  
National Naval Medical Center  
Bethesda, MD 20814

Director, Infectious Diseases Program Center  
Naval Medical Research Institute  
National Naval Medical Center  
Bethesda, MD 20814

Commander  
Chemical and Biological Sciences Division  
Army Research Office, P. O. Box 12211  
Research Triangle Park, NC 27709

Commander  
U.S. Army Research and Development Command  
Attn: SGRD-PLA  
Fort Detrick  
Frederick, MD 21701

Commander  
USAMRIID  
Fort Detrick  
Frederick, MD 21701

Directorate of Life Sciences  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, DC 20332

Mr. Robin A. Simpson  
Representative - Stanford; Administrative Contracting Officer  
Office of Naval Research Resident  
Room 165, Durand Building  
Stanford, CA 94305

**Final and Technical Reports Only**

Director, Naval Research Laboratory (6 copies)  
Attn: Technical Information Division, Code 2627  
Washington, DC

## DISTRIBUTION LIST

### Immunological Defense Program

#### Annual, Final and Technical Reports (one copy each except as noted)

Fritz H. Bach, M.D.  
Director, Immunology Research Center  
University of Minnesota  
Box 724, Mayo Memorial Bldg.  
420 Delaware St., SE  
Minneapolis, MN 55455

Francis A. Ennis, M.D.  
Department of Medicine  
University of Massachusetts  
Medical School  
55 Lake Avenue  
Worcester, MA 01605

Fred D. Finkelman, M.D.  
Department of Medicine  
Uniformed Services University  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814

Dr. Matthew J. Kluger  
Department of Physiology  
University of Michigan Med. School  
7620 Medical Science II Bldg.  
Ann Arbor, MI 48109

Dr. Vijaya Manohar  
Borrison Laboratories, Inc.  
5050 Beech Place  
Temple Hills, MD 20748

Dr. Ernest D. Marquez  
Bioassay Systems Corporation  
225 Wildwood Avenue  
Woburn, MA 01801

James J. Mond, M.D.  
Department of Medicine  
Uniformed Services University  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814

Dr. Donna S. Sieckmann  
Infectious Diseases Program Center  
Naval Medical Research Inst.  
National Naval Medical Center  
Bethesda, MD 20814

Dr. John D. Clements  
Department of Microbiology  
and Immunology  
Tulane University Medical Ctr.  
1430 Tulane Avenue  
New Orleans, LA 70112

Dr. Edward A. Havell  
Trudeau Institute  
P.O. Box 59  
Saranac Lake, NY 12983

Dr. Arthur G. Johnson  
Department of Medical  
Microbiology and Immunology  
University of Minnesota  
School of Medicine  
2205 East 5th Street  
Duluth, MN 55812

Dr. Philip Lake  
Immunologic Oncology Division  
Lombardy Cancer Center  
Georgetown Univ. School of Med  
Washington, DC 20007

W. John Martin, M.D., Ph.D.  
Laboratory, Dept. of Medicine  
Naval Hospital  
National Naval Medical Center  
Bethesda, MD 20814

Dr. Robert I. Mishell  
Dept. of Microbiology &  
Immunology  
Univ. of California, Berkeley  
Berkeley, CA 94720

Dr. Page S. Morahan  
Department of Microbiology  
Medical College of Pennsylvania  
3300 Henry Avenue  
Philadelphia, PA 19129

Dr. Alan L. Schmaljohn  
Department of Microbiology  
University of Maryland  
School of Medicine  
660 W Redwood Street  
Baltimore, MD 21201

Annual, Final and Technical Reports (Cont.)

David A. Stevens, M.D.  
Department of Medicine  
Santa Clara Valley Medical Center  
Stanford University  
751 S. Bascom Avenue  
San Jose, CA 95128

Dr. Barnet M. Sultzer  
Department of Microbiology &  
Immunology  
Downstate Medical Center  
450 Clarkson Avenue  
Brooklyn, NY 11203

Dr. Alvin L. Winters  
Department of Microbiology  
University of Alabama  
University, AL 35486

Dr. Phyllis R. Strauss  
Department of Biology  
Northeastern University  
360 Huntington Avenue  
Boston, MA 02115

G. Jeanette Thorbecke, M.D.  
Department of Pathology  
New York University  
School of Medicine  
550 First Avenue  
New York, NY 10016

Lyn Yaffe, M.D.  
Research Support Center  
Naval Medical Research Inst.  
National Naval Medical Center  
Bethesda, MD 20814

**END**

**FILMED**

1-85

**DTIC**